



WESTERN

JUNE 2018

GRDC™ GROWNOTES™



GRDC™

GRAINS RESEARCH
& DEVELOPMENT
CORPORATION

VETCH

SECTION 2

PLANNING

KEY POINTS | VARIETY CHOICE | AUSTRALIAN NATIONAL VETCH BREEDING PROGRAM | PLACE IN ROTATION – CONSIDERATIONS FOR FUTURE CROPS | VETCH BENEFITS TO CEREAL ROTATIONS | PADDOCK SELECTION – CONSIDERATIONS FOR A VETCH CROP

Planning

Key points

- Select varieties based on vetch species paddock situation and end use.
- Vetch should not be grown more than once within a five-year period in the same paddock; however, farmers sowing soft-seeded varieties grow vetch successfully every four years.
- Vetch is a break crop for foliar and root disease in cereals and oilseeds but could carry over some soilborne disease, such as *Sclerotinia*.
- Diseases of vetch can be transferred via soil, stubble or volunteers of other pulse crops.
- Residues of some herbicides (such as sulfonylureas) can reduce vetch growth and yield severely.

MORE INFORMATION

A paper on 'Pulse performance and agronomy update' was delivered by SARDI's Larn McMurray at the southern region GRDC Updates held in February 2017. <https://grdc.com.au/resources-and-publications/grdc-update-papers>

A paper on 'Image analysis to quantify pulse quality traits' was delivered by PhD student Linda McDonald at the southern region GRDC Updates held in February 2017. <https://grdc.com.au/resources-and-publications/grdc-update-papers>

IN FOCUS

Australian National Vetch Breeding Program

The Australian National Vetch Breeding Program (ANVBP) collaborates with growers, scientists and agronomists in South Australia, Victoria, New South Wales, Western Australia and Tasmania, running trials aimed at identifying the best-performing varieties and end uses for vetch in particular areas.

There are also companies that have released vetch varieties in Australia.

ANVBP focuses on breeding varieties with:

- high yields of grain and dry matter
- resistance to rust, Ascochyta blight and Botrytis grey mould
- soft seed to avoid volunteer vetch weed problems in following crops
- lower toxins in the grain so it is suitable as a stock feed
- varieties adapted to lower-rainfall areas where other pasture legume or pulse crops are performing poorly
- non-shattering pods.

Species and variety choice is determined by the paddock situation and end use. The place in rotation is determined by end use, weed burden, herbicide residues and disease carryover.

MORE INFORMATION

South Australian Sowing Guide 2017, <https://grdc.com.au/resources-and-publications/all-publications/publications/2016/12/sa-sowingguide2017>

Vetch seed suppliers

Auswest Seeds, <http://www.auswestseeds.com.au/vetch1>

Heritage Seeds, <http://www.heritage-seeds.com.au/forage-pasture-3/clover-pasture-legumes/vetch>

Pasture Genetics, <http://pasturegenetics.com>

For farm trial information, please see: Emerging forage legume varieties (2013) <http://www.farmtrials.com.au/trial/14135>

MORE INFORMATION

For a paper on one of the latest trial results on vetch please see <http://www.giwa.org.au/2017researchupdates>

‘Love them legumes – recent adventures with lentils, peas and beans’, Update paper by Mark Seymour – www.giwa.org.au/_literature_225530/S15_Mark_Seymour_2017

A concise overview of this presentation by Mark Seymour, Department of Primary Industries and Regional Development, can be viewed at: <https://youtu.be/0wXEHsyeflA>

2.1 Variety choice

Vetch varieties can be selected from three species that have different end uses. The first step in selecting a vetch variety is to consider the likely rainfall, the second is to determine end use.

Different species have different end uses:¹

- Common or grain vetch – *Vicia sativa* subspecies *sativa* – forage, manure crop, grain for stockfeed, seed.
- Purple vetch – *Vicia benghalensis* subspecies *benghalensis* – forage and manure crop, seed.
- Woolly pod vetch – *Vicia villosa* subspecies *dasycarpa* and *eriocarpa* – forage and manure, seed.

Some Common vetch varieties, e.g. Morava[®], Rasina[®], Timok[®] and Volga[®], produce high yields of forage and grain.

Vetch species and varieties differ in their adaptation to rainfall regions, in their end use suitability, productivity and percentage of hard seed. When selecting a vetch variety the following factors all need to be considered (see also [Table 1](#) and [Table 2](#)):

- rainfall
- suitability for early grazing
- forage and/or grain production
- suitability for stock feed
- percent hard seed
- maturity – flowering in relation to frost and heat stress
- pod shatter – if for grain
- level of anti-nutritional factors – if for stock feed
- disease resistance – in-crop control of foliar diseases can be expensive
- herbicide tolerance – impact depends on the herbicide, soil type and rainfall and time since application (see [Section 8](#))

If vetch is sown in a mixture with cereals for cutting for silage or hay, a cereal variety with similar maturity should be selected in order to maximise quality at cutting.

In trials in WA in 2016 five new vetch lines were trialled as well as Timok[®], Morava[®] and Volga[®]. In a year with a cool wet winter and cool spring the vetch lines yielded between 1.9 to 2.2t/ha and the varieties all produced about 1.8t/ha. This suggests vetch grain yields in WA may be lower than those suggested in [Table 1](#).

Species and varieties have similar susceptibility to insect pests (see [Section 2.5.5](#)). From early growth, through to pod maturity, they are susceptible to bluegreen and cowpea aphids, as well as to native budworm during pod formation and filling.

The lists and tables present varieties in alphabetical order. [Section 2.1.1](#) to [Section 2.1.3](#) indicate if an improved or replacement variety is available.

¹ PIRSA/GRDC (2010) Vetch: The Ute Guide. PIRSA/GRDC – <https://grdc.com.au/vetch-the-ute-guide>

SECTION 2 VETCH

FEEDBACK



Photo 1: Not all vetch species are the same, nor can they all be grown for the same outputs. Variety choice must match vetch to the growing environment and end use.

SECTION 2 VETCH

FEEDBACK

Table 1: Adaptation, usage and production of varieties of vetch in the main subspecies grown in Australia based on seven trials in South Australia, 2013–15.

Variety	Rainfall (mm)	Suitability for early grazing	Forage production	DM (t/h)	Grain production	Grain (t/ha)	Grain in stockfeed	% hard seed
Common or Grain vetch – <i>Vicia sativa ssp. sativa</i>								
Blanchefleur	350–450 g <350–400 f	Good	Moderate	4.03	Good	2.15	Yes Max for pigs 20%	5–10
Cummins	<350–450 g <350–400 f	Moderate	Moderate	–	Good	–	Yes Max for pigs 20%	5–15
Languedoc	<350–400 g <350 f	Good	Moderate	–	Good	–	Yes Max for pigs 20%	5–15
Morava [Ⓛ]	400–>600 g <350–>600 f	Poor	Good	5.06	Good	2.16	Yes Max for pigs 25%	0
Rasina [Ⓛ]	350–600 g <350–450 f	Good	Moderate	4.7	Good	2.37	Yes Max for pigs 25%	0
Timok [Ⓛ]	<350–>600 g <350–>600 f	Good	V. good	5.26	Good	2.48	Yes	0–2
Volga [Ⓛ]	<350–450 g <350–450 f	Moderate	Good	5.51	Very good	2.75	Yes Max for pigs 25%	2–5
Purple vetch – <i>Vicia villosa ssp. benghalensis</i>								
Benatas ¹	350–800 f	Moderate	Good	9.71	Poor	–	No	Low
Popany	400–>600 f	Poor	Good	5.28	Poor	–	No	5–10
Woolly pod vetch – <i>Vicia villosa ssp. dasycarpa and eriocarpa</i>								
Capello [Ⓛ]	400–>600 f	Poor ²	Very good	6.23	Poor	–	No	15–20
Haymaker [Ⓛ]	400–>600 f	Poor ²	Very good	6.26	Poor	–	No	20–30
Namoi	400–>600 f	Very poor ²	Very good	–	Poor	–	No	>80
RM4 [Ⓛ]	<350–>600 f	Moderate ²	Very good	6.71	Moderate	–	No	2–5

Notes: 1. Source: Tasglobal Seeds; 2. Graze post 10th node stage to end of flower.

Key: g= for grain, f = for fodder

DM assessments made at flowering.

Source: SARDI (2013, 2016, 2017) South Australian Sowing Guide. SARDI <https://grdc.com.au/SA-SowingGuide2017>

SECTION 2 VETCH

FEEDBACK

Table 2: Additional characteristics that influence variety choice.

Characteristic	Maturity	Shattering	GBC ¹	Disease rating		
				Ascochyta blight	Botrytis	Rust
Common vetch – <i>Vicia sativa</i> ssp. <i>sativa</i>						
Blanchefleur	Early	MS	0.9–1.2%	MS	S	VS
Cummins	Early-mid	MS	1.2%	MS	S	VS
Languedoc	Very early	MS	1.0–1.6%	MR	S	VS
Morava [Ⓛ]	Late	R	0.65%	S	VS	R
Rasina [Ⓛ]	Early	MR	0.66–0.85%	MS	S	R
Timok [Ⓛ]	Mid	MR	0.57%	MS	S	R
Volga [Ⓛ]	Early	MR	0.54%	MS	S	R
Purple vetch – <i>Vicia villosa</i> ssp. <i>benghalensis</i>						
Benatas	Late	–	–	–	–	–
Popany	Very late	MR		MS	VS	R
Woolly pod vetch – <i>Vicia villosa</i> ssp. <i>dasycarpa</i> and <i>eriocarpa</i>						
Capello [Ⓛ]	Late	R		MR	VS	R
Haymaker [Ⓛ]	Late	R		MR	VS	R
Namoi	Very late	R		MR	VS	R
RM4 [Ⓛ]	Mid	MR		MR	VS	R

Note: 1. GBC = gamma-glutamyl-beta-cyanoalanine, an anti-nutritional factor.

Key: R = resistant, MR = moderately resistant, MS = moderately susceptible, S = susceptible, VS = very susceptible

Source: R Matic (2016) personal communication. Australian National Vetch Breeding Program



Photo 2: Days to full flowering from sowing by variety, sown on the same day.

Photo: Wayne Hawthorne

2.1.1 Common or Grain vetch (*Vicia sativa* ssp. *sativa*)

The newer varieties of Common vetch – Morava[Ⓛ], Rasina[Ⓛ], Timok[Ⓛ] and Volga[Ⓛ] – have improved rust resistance, productivity and soft seed percentage. The following list is in alphabetical order and indicates if an improved or replacement variety is available.

Blanchefleur

Description: hairy, oblong, squarish adult leaves with medium-green foliage. White flowers on short stalks. Long, narrow pods containing reddish-brown, mottled seed. Pillow-shaped seed, which is orange when split. Seed size: 6.6 g/100 seeds.

The market for varieties with orange cotyledons, like Blanchefleur, is limited to domestic markets for birdseed and sowing seed for grazing and manure crops. This restriction is to prevent the substitution of vetch grain for lentil. Both vetch and lentil are on the Australian Government's prescribed grain list.²

Prior to the release of Morava[Ⓛ], Blanchefleur was the preferred grain variety in areas of >350 mm rainfall. Blanchefleur has mid-maturity and is well suited to medium to high-rainfall areas where rust is not a regular problem as it is very susceptible to rust.

Replaced by Rasina[Ⓛ] in low-rainfall areas and Morava[Ⓛ] in higher-rainfall areas.

Cummins

This is a mid to early maturing, white-flowering variety selected from Languedoc. Seed size: 6.1 g/100 seeds.

It is well adapted to medium to low-rainfall areas where it generally yields higher than Blanchefleur. Cummins is susceptible to rust and moderately susceptible to Ascochyta blight.

Replaced by Rasina[Ⓛ] in low-rainfall areas and Morava[Ⓛ] in higher-rainfall areas.

Languedoc

Description: hairy, oblong, squarish adult leaves with medium-green foliage. Light-purple flowers on short stalks. Long, narrow pods containing brown-grey, pillow-shaped seed, which is beige when split. Seed size: 6.7 g/100 seeds.

It is an early-flowering and maturing variety recommended for low-rainfall areas, although it can lodge severely if there is heavy rainfall when it is ripe, making harvest difficult. Languedoc is generally higher yielding than Blanchefleur in areas with <350 mm rainfall. Its hard seed content is generally around 5–10% and it is highly susceptible to rust.

Replaced by Rasina[Ⓛ].

Morava[Ⓛ]

Description: hairy, oblong, large adult leaves with dark-green foliage. Dark-purple flowers on short stalks. Very long, narrow pods containing large, dark-brown, pillow-shaped seed, which is beige when split. Seed size: 8.3 g/100 seeds.

Developed in 1998 by the Australian National Vetch Breeding Program (ANVBP), Morava[Ⓛ] is a late-flowering vetch variety with 100% soft seeds. It has large seed and is more resistant to shattering than other vetch varieties.

Morava[Ⓛ] has superior grain yield to other vetches in the high-rainfall areas and, in all other areas, has a higher yield than Blanchefleur, Languedoc and Cummins where rust is present.

² Australian Government Department of Agriculture and Water Resources (2016) <http://www.agriculture.gov.au/export/controlled-goods/plants-plant-products/ian/03/08> Advice Notice no. 2003/08: Export of Split Vetch Prohibited

FEEDBACK

It has a lower level of the anti-nutritional factor gamma-glutamyl-beta-cyanoalanine (GBC) (0.65%) than Blanchefleur and Languedoc. Morava[®] produces high herbage yields. Morava[®] is later flowering and maturing than Blanchefleur and grain yield is reduced in seasons with dry finishes.

Morava[®] is rust resistant, susceptible to Ascochyta blight and very susceptible to Botrytis grey mould, because it produces very high biomass in wetter areas.

Morava[®] is a PBR variety and seed can be sourced from Heritage Seeds.

Rasina[®]

Description: hairy, oblong, squarish adult leaves with medium-green foliage. Light-purple flowers on short stalks. Long, narrow pods containing seeds with a dark-brown, speckled coat; pillow-shaped seed is dark beige-greenish when split. Seed size: 6.9 g/100 seeds.

Developed in 2006 by ANVBP, Rasina[®] is an earlier-flowering, soft-seeded variety that replaces Languedoc, Blanchefleur and Cummins in low to medium-rainfall areas for grain production. It is a soft-seeded variety and has a low level of the anti-nutritional factor GBC (0.6–0.8%).

A significant advantage over Languedoc, Blanchefleur and Cummins is its resistance to rust. Rasina[®] is not expected to replace Morava[®] in higher-rainfall districts or for hay production.

Rasina[®] is a PBR variety and seed can be sourced from Heritage Seeds.

Timok[®]

Description: dark-green leaves, which are convex to straight in early stage. Light-violet flowers. Pods medium to long and width medium to wide. Seed coat brown with black speckling, with grey-brown cotyledons. Seed size: 6.9 g/100 seeds.

Released in 2013 by ANVBP, Timok[®] was bred to complement Morava[®] in mid to high-rainfall areas for grain and especially for silage and hay production. It is a soft-seeded variety and has a low level of the anti-nutritional factor GBC (0.57%).

Timok[®] has better early growth than Morava[®] and will improve the reliability of vetch and economic production in cropping systems especially in mid-rainfall areas (350–450 mm per year). Morava[®] is still the preferable variety for hay and silage in areas with >450 mm rainfall per year.

It is a high-yielding and highly rust-resistant Common vetch variety, which is moderately susceptible to Ascochyta blight and susceptible to Botrytis grey mould.

Timok[®] is a PBR variety and seed can be sourced from Pasture Genetics.

Volga[®]

Description: leaves are concave and medium green to dark green. Flowers are medium-violet and pods medium to long. Seed coat is brown with blue-black speckling, cotyledons grey brown. Seed size: 7.8 g/100 seeds.

Released in 2013 by ANVBP, Volga[®] is a high-yielding grain and herbage variety for low and mid-rainfall areas. Its early flowering makes it particularly suited to shorter-season areas where the growing season finishes sharply.

It has a small proportion of hard seed and a low level of the anti-nutritional factor GBC (0.54%).

Volga[®] has good initial establishment and early maturity. It is rust resistant and moderately susceptible to Ascochyta blight. It is earlier flowering and maturing than Blanchefleur and Rasina[®].

Volga[®] is a PBR variety and seed can be sourced from Heritage Seeds.

2.1.2 Purple vetch (*Vicia benghalensis ssp. benghalensis*)

Benetas

Description: similar to Popany, but no detailed description available.

Developed by Tasglobal Seeds, Benetas produces high forage yields with good early spring vigour. Benetas is later flowering than Popany and has improved cold tolerance during vegetative growth (has survived to -7°C). It is also tolerant of moderate waterlogging. These characteristics make it especially suited to cooler, higher-rainfall regions.

No disease-resistance data is currently available.

Benetas seed can be sourced from AusWest Seeds and Ardent Seeds in Tasmania.

Popany

Description: oblong, narrow leaves with medium to dark-green foliage. Purple flowers with dark lips on stalks the length of the leaf. Medium-length brown pods containing velvety black, globular seed with a white hilum, which is yellow when split. Seed size: 4.5 g/100 seeds. A small proportion of seed is hard.

Popany is a late-maturing variety good for hay and silage production in mid to high-rainfall areas.

Grain yield is significantly lower than yields of Common vetch varieties and seed size is smaller. This variety is resistant to rust but susceptible to Ascochyta blight and chocolate spot.

Popany seed can be sourced from farmer sale as well as several seed houses.

2.1.3 Woolly pod vetch

All currently available Woolly pod vetch varieties can be grazed from 10 nodes to podset because of anti-nutritional issues outside these growth stages.

Unlike other vetch species they are all moderately resistant to Ascochyta blight. If disease conditions occur these varieties are susceptible to Botrytis grey mould.

Grain can only be sold as seed.

Capello[Ⓛ] and Haymaker[Ⓛ] (*Vicia villosa ssp. dasycarpa*)

Description: oblong, narrow leaves with medium-green foliage. Purple flowers, with pink inner, on stalks longer than the leaf. Short, beaked pods containing dark-brown, globular seed, which is bright yellow when split. Seed size: 4.5 g/100 seeds. Moderately hard-seeded.

Haymaker[Ⓛ] and Capello[Ⓛ] are selected softseed varieties from Namoi. They are lower in grain yield but much higher in dry matter production than Common vetch varieties in rainfall areas of >450 mm per year. These two varieties are very good for hay and silage production in areas where there is >400 mm of annual rainfall.

Both varieties are owned by Heritage Seeds.

Namoi

Description: oblong, narrow leaves with medium-green foliage. Purple flowers, with pink inner, on stalks longer than the leaf. Short, beaked pods containing dark-brown, globular seed, which is bright yellow when split. Seed size: 4.5 g/100 seeds. Very hard-seeded.

Namoi can be sourced from farmer sale as well as several seed houses.

RM4[®] (*Vicia villosa* ssp. *eriocarpa*)

Description: oblong, narrow leaves with medium-green foliage. Purple flowers, with pink inner, on stalks longer than the leaf. Short, beaked pods containing dark-brown, globular seed, which is bright yellow when split. Seed size: 4.5 g/100 seeds. Soft-seeded.

Bred by the ANVBP and released in 2014, RM4[®] is a multipurpose variety that can be used for silage or hay, grazing, a manure crop or for seed.

RM4[®] has moderate early growth, better than other Woolly pod varieties. It produces more dry matter than Capello[®] and Haymaker[®] in low and mid-rainfall areas and is also suitable for higher-rainfall areas (>400–650 mm per year). Its early maturity helps RM4[®] produce more dry matter than other Woolly pod varieties when the growing season finishes sharply. It is excellent for improving soil nitrogen and soil structure.

RM4[®] is a PBR variety and can be sourced from Heritage Seeds.

2.2 Australian National Vetch Breeding Program

The Australian National Vetch Breeding Program (ANVBP) collaborates with growers, scientists and agronomists in South Australia, Victoria, New South Wales, Western Australia and Tasmania, running trials aimed at identifying the best-performing varieties and end uses for vetch in particular areas.

The ANVBP allows producers and end users in different regions to observe how present varieties are performing and to evaluate potential new varieties that could be suited to those areas. ANVBP focuses on breeding varieties with:

- high yields of grain and dry matter
- resistance to rust, Ascochyta blight and Botrytis grey mould
- soft seed to avoid volunteer vetch weed problems in following crops
- lower toxins in the grain so it is suitable as stock feed
- varieties adapted to lower-rainfall areas where other pasture legume or pulse crops are performing poorly
- non-shattering pods.

There are also a few private breeding companies producing vetch for the Australian market.

FEEDBACK

i MORE INFORMATION

Information about PreDicta B,
http://pir.sa.gov.au/research/services/molecular_diagnostics/predicta_b

2.3 Place in rotation – considerations for future crops

To reduce risk of disease and crop contamination, ideally vetch should not be grown more than once within five years in the same paddock and should not be sown adjacent to vetch, bean or lentil stubbles. In reality, a one-in-four-year rotation has been found to be successful, especially when sowing soft-seed varieties.

2.3.1 Disease

Vetches provide a good break crop for root and some foliar diseases of cereals and canola. Vetch can host root-lesion and stem nematode, as well as several root diseases that can affect cereals and other pulse crops (see Table 3). Carryover of some soil and stubble-borne diseases can be tested using the PreDicta B DNA-based testing service; the tests relevant to vetches measure soil inoculum levels of *Rhizoctonia* bare patch (see Figure 1) and stem nematode.

There can be some disease crossover between vetch and faba bean, lentil, field pea, chickpea and lupin (see Table 3). Disease crossover can come from adjacent crops, from stubble or from the soil, depending on the disease (see Section 3.5 Carryover diseases).

Table 3: Vetch diseases and potential for cross-infection from other pulses.

Disease	Vetch	Lentil	Field pea	Faba bean	Chickpea	Lupin
Ascochyta blight# (<i>Ascochyta fabae</i>)	*	^	^	**	^	^
Botrytis grey mould (<i>Botrytis cinerea</i>)	**	**	*	**	**	*
Chocolate spot (<i>Botrytis fabae</i>)	**	**	^	**	^	^
Rust (<i>Uromyces viciae-fabae</i>)	**	^	*	**	^	^
<i>Sclerotinia</i> stem rot# (<i>Sclerotinia</i> sp.)	**	**	*	**	**	**
Stem nematode (<i>Ditylenchus dipsaci</i>)	*	*	**	*	*	^
Viruses: non-persistent AMV, BBWV, BYMV, CYVV and PSbMV	**	**	**	**	**	**
Viruses: persistent BLRV, BWYV, SbDV and SCSV	**	**	**	*	**	**
Root rots <i>Fusarium</i> sp.	*	*	*	*	*	*
<i>Phoma</i> sp.	*	*	**	*	**	*
<i>Pythium</i> sp.	*	*	*	*	*	*
<i>Rhizoctonia</i> sp.	**	**	**	**	**	**

Key: * This disease occurs on this crop but has not caused major damage ** This disease has caused major damage on this crop
 ^ Non-host # Species differ between crops

Source: PIRSA/GRDC (2010) Vetch: The Ute Guide. PIRSA/GRDC – <https://grdc.com.au/vetch-the-ute-guide>

FEEDBACK




Figure 1: The effect of crop rotation, including vetch, pea and lupin, on *Rhizoctonia solani* AG8 inoculum level at low-rainfall sites in Victoria, 2011. As few as five ryegrass plants/m² can maintain or increase *Rhizoctonia* levels in a break crop.

Source: GRDC (2012) *GroundCover™* Supplement 98, <https://grdc.com.au/resources-and-publications/groundcover/ground-cover-issue-98-may-june-2012>

i MORE INFORMATION

WeedSmart,
<https://weedsmart.org.au>

Australian Pesticides and Veterinary
Medicines Authority, apvma.gov.au

2.3.2 Weeds

Self-sown vetches from hard-seeded varieties can cause admixture quality problems in some crops at harvest. Separating vetch from field pea and lentil in the harvesting process can be difficult. Rotations should be designed to avoid unwanted contamination of vetch in these crops.

The control of vetch in other pulse crops still largely relies on pre-season seedbank management.

WeedSmart offers a suite of tools to help plan weed management for future crops on a paddock-by-paddock basis.

In trials in the Mallee (2011–13) to identify profitable rotations for brome grass control, the best brome grass control and gross margins were achieved with a sequence of vetch and Clearfield® canola or Clearfield® wheat followed by wheat.³

³ GRDC & BCG (2013) Mallee rotations that beat brome grass and make money. <http://www.farmtrials.com.au/trial/16654>

SECTION 2 VETCH

FEEDBACK




Figure 2: Cumulative gross margin (\$/ha) and final brome grass populations on 29 August 2013 (plants/m²) in the Mallee (2011–13). 2011–13 GM (\$/ha): $P < 0.001$, $LSD = \$238$, $CV = 20\%$; final brome (plants/m²): $P < 0.001$, $LSD = 13$ plants/m², $CV = 18\%$.

Source: GRDC & BCG (2013) Mallee rotations that beat brome grass and make money. GRDC <http://www.farmtrials.com.au/trial/16654>

Where vetch exists as a volunteer in cereals there are several herbicide options, including mixes of phenoxy-based herbicides, such as MCPA that can be used depending in which crop the volunteers are present. Check herbicide labels for registration for use in vetch on the APVMA website.



Photo 3: (Left) tares on the headland of a vetch crop. The control of vetch or tares in pulse crops largely relies on pre-season weed seedbank management. (Right) tares (pictured on the left) is a smaller, more spindly plant than modern grain vetch plants.

Photo: Emma Leonard, AgriKnowHow

2.3.3 Nitrogen fixation

Results from the ANVBP across five sites over three years have shown increases in soil nitrogen after vetch was grown for grain (56 kg/ha), hay (94 kg/ha) and green manure (154 kg/ha). (See Section 1, [Table 2](#).)

2.3.4 Stubble cover

Vetch stubble residues lack bulk and provide only partial protection to the soil after harvest. Vetch stubble should either not be grazed or grazed with caution to ensure adequate stubble cover is maintained to minimise risk of wind and water erosion on sandy soils and sloping paddocks. Vetch holds the soil better than field pea as it produces more cover and surface roots.

At seeding, the stubble of unharvested crops can be a problem due to the long vines leading to soil build-up under the seeder. This stubble may have to be harrowed or disced ahead of seeding to reduce this problem.

2.3.5 Soil moisture reserves

In crop-sequencing trials, vetch consistently fixed more nitrogen and used more water than pea, probably because of its longer growing season and greater dry matter accumulation. However, an early manure crop or cutting as silage results in more water remaining in the profile for the following crop than if vetch was grown to maturity and harvested as grain.⁴

2.4 Vetch benefits to cereal rotations

- Increased yields of following cereal crops.
- Allows an extended phase of cropping.
- Decreases many cereal diseases – grass-free vetch crops break the life cycle of root diseases, crown rot, take-all and *Rhizoctonia*.
- Controls grass weeds – cutting for forage, using grass-selective herbicides or manuring can be used with vetches to control weeds, such as brome grass and barley grass, which are difficult to control in some other crops.
- Allows for crop-topping to prevent herbicide-resistant weeds from setting seed.
- Available soil nitrogen is improved.
- Well adapted to no-till, standing-stubble systems aimed at improving soil structure and fertility.⁵

2.5 Paddock selection – considerations for a vetch crop

2.5.1 Soil type

Vetch will grow on a wide range of soil types from light sandy soils to heavier clay soils.

On light, sandy soils the Common and Woolly pod varieties perform well. All perform well on loam clay soils but the best production comes from soils with good fertility.

Vetch prefers alkaline soils (pH 5.2–8.2) but the variety Benatas, for example, has been found to perform well on slightly more acidic soils.⁶

Common vetch will not survive prolonged waterlogging. The vetch subspecies Purple vetch and Woolly pod will tolerate waterlogging and survive better than other crops, such as oats.

Vetch is less tolerant of acidity than lupin

Vetch is moderately sensitive to salinity, and can have difficulty accessing water and nutrients from saline layers in the soil. Soil chloride levels >600 mg/kg have been found to reduce root growth in crops such as chickpea, lentil and linseed.

Vetch is classified among the 'medium' group for sensitivity of all field crops to sodic soil conditions.

MORE INFORMATION

GroundCover™ TV, Break Crops Decision Tool,
https://www.youtube.com/watch?v=Zq_jnpLxGDw&feature=youtu.be_gdata_player

4 M Peoples (2012) *GroundCover™ Supplement 98*. (2012) GRDC <https://grdc.com.au/resources-and-publications/groundcover/ground-cover-issue-98-may-june-2012>

5 PIRSA/GRDC (2010) *Vetch: The Ute Guide*. PIRSA/GRDC – <https://grdc.com.au/vetch-the-ute-guide>

6 C Gazey, J Carson (2016) *Managing Soil Acidity – WA. Fact Sheets*, Department of Agriculture and Food, Western Australia, <http://www.soilquality.org.au/factsheets/managing-soil-acidity-western-australia>

 MORE INFORMATION

GRDC Podcast, Herbicide residues, radio <https://grdc.com.au/Media-Centre/GRDC-Podcasts/Driving-Agronomy-Podcasts/2016/05/Herbicide-Residues>

2.5.2 Herbicide residue

Rotations must also take into account herbicide residues and plant-back requirements in relation to soil pH, texture and organic matter content, especially after drought conditions. Herbicide residue impacts are more pressing where rainfall has been minimal and, in many cases, where the soil type is heavier.

It is very important to know the chemical history of the paddock for at least two seasons, this includes knowledge of:

- the chemical used
- the group to which the chemical belongs
- the plant-back periods
- the soil pH (which affects the half-life and longevity of herbicides)
- rainfall
- other requirements for specific herbicide breakdown (check labels).

Residues of sulfonylureas (SUs) (chlorpyralid) and imidazolinones (IMIs) (Groups B and I) can be particularly harmful in alkaline soils. Always check withholding periods. For example, there is a 10-month withholding period for sowing vetch after the use of the Group B herbicide active ingredient imazethapyr.

2.5.3 Sowing into cereal stubble

Vetches grow well when sown after cereal or oilseed crops. Paddocks with adequate standing stubble cover provide a trellis, lifting the crop off the soil surface, which is good for grain crops.

If sowing for hay crops, stubble cover is also good to provide early protection but stubble should be rolled to the surface to minimise incorporation with the cut hay.

2.5.4 Disease

Controlling foliar diseases in vetch can be expensive. Effective disease management relies on variety and paddock selection, plus the use of clean seed, best agronomic practice and the strategic use of fungicides (see [Section 7.1 Disease management](#)).

Time of sowing and seasonal conditions influence the incidence of foliar disease, particularly Botrytis grey mould (BGM). A dense canopy is conducive to BGM in a wet season.

See [Table 3](#) for the potential disease carryover from other crops in the rotation. A four to five-year break from vetch and disease carryover crops is required to minimise disease transfer and vetch should not be sown adjacent to vetch, bean or lentil stubbles.

2.5.5 Insect pests

Vetch varieties show little difference in their pest susceptibility. Generally, damage from pests of emerging crops, such as snails, slugs, millipedes and earwigs, is not a major problem. This is partly because in vetch, **unlike lupin**, cotyledons do not grow above the soil surface, so plants can reshoot underground if the tops have been eaten off.

All current vetch varieties are susceptible in early growth stages to redlegged earth mite and lucerne flea. Most are susceptible to bluegreen and cowpea aphids from early growth through to pod maturity, as well as to native budworm during pod formation and filling (see [Section 6.1 Pest management](#)).

2.5.6 Cross-pollination

If Woolly pod vetch crops are to be used as seed, they should be planted more than 400 m from other varieties to reduce the risk of cross-pollination. Common vetch varieties are self-pollinated and do not require a distance of more than 50 m from the other vetches.