



WESTERN
SEPTEMBER 2018

GRDC™ **GROWNOTES™**



GRDC™
GRAINS RESEARCH
& DEVELOPMENT
CORPORATION

CHICKPEA

CONTENTS

Contents

Introduction

Key messagesxxiv

A.1 Crop overviewxxiv

Desixxv

Kabulixxvi

A.1.1 Pulsesxxvi

A.1.2 Quality attributesxxvii

A.1.3 Nutritional informationxxviii

A.1.4 Agronomy at a glancexxviii

A.2 Brief historyxxix

A.3 GRDC chickpea breeding investmentxxx

A.4 Keywordsxxxi

1 Planning/Paddock preparation

Key messages1

1.1 Paddock selection 2

Key points2

1.1.1 Avoid deep gilgai or heavily contoured country3

1.1.2 Soil3

Adaptation and seed yield of cool-season grain legumes in

Mediterranean environments of south-western Australia4

Subsoil5

1.1.3 Soil pH5

Managing soil pH6

1.1.4 Rainfall6

1.1.5 Physical constraints7

1.1.6 Nutrient constraints7

1.1.7 Biological constraints7

1.1.8 Problematic paddocks8

1.1.9 Stubble retention8

Stubble and its impact on temperature in chickpea crops10

Conclusions11

Bunching and clumping of stubble11

1.1.10 Soil testing guide12

Accurate results13

Taking the test13

When to collect samples13

	<i>Regular tests build better profile.....</i>	13
	<i>Selecting your samples.....</i>	13
	<i>Account for variability.....</i>	14
	<i>Sampling sites.....</i>	14
	<i>Handle with care.....</i>	14
	<i>Interpreting the results.....</i>	14
	<i>How to take a soil sample.....</i>	14
1.2	Paddock rotation and history.....	16
1.2.1	Break cropping.....	16
1.2.2	Chickpea as a rotation crop.....	16
1.2.3	Pulse effects on cereal yield.....	17
	<i>Nitrate – N benefit for following cereals.....</i>	18
	<i>Crown rot.....</i>	19
1.2.4	Understanding soils and pulse crop constraints.....	20
1.3	Fallow weed control.....	22
1.3.1	Management strategies.....	24
	<i>Herbicide application.....</i>	24
	<i>Double-knock strategies.....</i>	25
	<i>Grazing summer weeds.....</i>	25
1.4	Fallow chemical plant-back effects.....	26
1.4.1	Herbicide residues in soil.....	27
	<i>Conditions required for breakdown.....</i>	29
1.5	Seedbed requirements.....	29
	<i>No-till.....</i>	29
	<i>Tillage.....</i>	29
1.6	Soil moisture.....	30
1.6.1	Dryland.....	30
	<i>Preserving dryland fertility.....</i>	30
1.6.2	Irrigation.....	31
	<i>Factors to consider when planning for irrigated chickpea production:.....</i>	32
	<i>Irrigation techniques to reduce the period of waterlogging:.....</i>	34
	<i>Spray irrigation.....</i>	34
	<i>Irrigation management strategy for chickpea.....</i>	34
	<i>High value kabuli chickpea production in the Ord River Irrigation Area.....</i>	35
1.7	Yield and targets.....	37
	<i>Critical period for chickpea yield.....</i>	38
	<i>Estimating crop yields.....</i>	40
	<i>Yield Prophet.....</i>	40
1.7.1	Seasonal outlook.....	41
	<i>Australian CliMate.....</i>	41
1.7.2	Fallow moisture.....	42
	<i>HowWet/N?.....</i>	42
1.7.3	Water Use Efficiency.....	43
	<i>Chickpea Water Use Efficiency in relation to cropping system,</i>	

	<i>cultivar, soil nitrogen and Rhizobial inoculation in semiarid environments</i>	44
	<i>Managing to optimise Water Use Efficiency</i>	45
1.7.4	Nitrogen use efficiency.....	47
1.7.5	Double crop options.....	47
	<i>Strategic double cropping on Vertisols: A viable rainfed cropping option in India to increase productivity and reduce risk</i>	48
1.8	Disease status of paddock	48
1.8.1	Cropping history effects.....	49
1.9	Nematode status of paddock	49
1.9.1	Effects of cropping history on nematode status.....	49
1.10	Testing soil for disease and nematodes	50
1.10.1	Soil and plant testing services for diagnosing root diseases.....	50
	<i>Confirmation of diagnosis</i>	51
1.11	Insect status of paddock	52
1.11.1	Insect sampling of soil	52
	<i>Insect ID: The Ute Guide</i>	53
1.11.2	Effect of cropping history.....	54
2	Pre-planting	1
	<i>Key messages</i>	1
2.1.1	Choosing a variety.....	1
	<i>Area of adaptation</i>	2
	<i>Evaluation of yield potential</i>	2
2.2	Varietal performance and ratings yield	3
2.3	Planting seed quality	5
	<i>Effect of poor quality seed on yield</i>	6
	<i>Grower-retained seed</i>	7
	<i>Grading</i>	8
	<i>Handling seed</i>	8
2.3.1	Testing for seed quality	9
2.3.2	Seed size.....	9
2.3.3	Seed germination and vigour	9
	<i>Testing seed vigour</i>	10
	<i>Weed contamination testing</i>	12
	<i>Disease testing</i>	12
2.3.4	Seed purity.....	12
2.3.5	Seed storage	14
	<i>Insect pests in storage</i>	15
2.3.6	Safe rates of fertiliser sown with the seed	15
2.4	Future breeding directions	16
3	Planting	1
	<i>Key messages</i>	1
3.1	Inoculation	1

CONTENTS CHICKPEAS

TABLE OF CONTENTS

FEEDBACK

	<i>Nutrient uptake and yield of chickpea (Cicer arietinum L.) inoculated with plant growth- promoting rhizobacteria</i>	3
3.1.1	Inoculation checklist	4
3.1.2	Inoculant types	4
	<i>Peat inoculum</i>	4
	<i>In-furrow water injection</i>	4
	<i>Granular forms</i>	4
3.1.3	Choosing an inoculant type	5
3.1.4	When to inoculate	6
3.1.5	Inoculum survival	8
3.1.6	Inoculant quality assurance	8
3.1.7	Inoculation methods	8
	<i>For chickpeas:</i>	9
3.1.8	Inoculum slurry	9
3.1.9	In-furrow water injection	10
3.1.10	Inoculant application trials	12
3.1.11	Compatibility with other major factors	13
	<i>Pesticides</i>	13
	<i>Fungicides</i>	13
	<i>Trace elements</i>	14
3.1.12	Nodulation and nitrogen fixation	14
	<i>Chickpea Water Use Efficiency in relation to cropping system, cultivar, soil nitrogen and Rhizobial inoculation in semiarid environments</i>	16
3.1.13	Monitoring nodulation	16
3.1.14	Use of nitrogen in inoculation	20
3.2	Seed treatments	20
3.3	Time of sowing	21
	<i>When to sow chickpea in south-western Australia</i>	23
3.3.1	Frost damage	24
	<i>Damage to vegetative growth:</i>	24
	<i>Damage to flowers and pods:</i>	24
3.4	Seed rate	25
	<i>Optimum plant density of desi chickpea (Cicer arietinum L.) increases with increasing yield potential in south-western Australia</i>	26
3.4.1	Calculating seed requirements/sowing rate	26
3.5	Targeted plant population and row spacing	27
3.5.1	Wide rows (50–100 cm) offer:	28
3.5.2	Narrow rows (15–40 cm) offer:	29
	<i>Yield response of kabuli and desi chickpea (Cicer arietinum L.) genotypes to row spacing in southern Australia</i>	30
3.6	Row placement	30
3.6.1	Row orientation	31
	<i>East-west crop orientation</i>	31
3.7	Sowing depth	32

	<i>Sowing Depth for Chickpea, Faba Bean and Lentil in a Mediterranean-type Environment of South-western Australia</i>	33
3.7.1	Deep seeding strategies	34
	<i>Chickpea seedling features</i>	35
	<i>Deep planting checklist</i>	35
	<i>Deep planting method</i>	36
3.8	Sowing equipment	37
3.8.1	Seeders.....	37
3.8.2	Airseeders.....	38
3.8.3	Seeder and tyne comparisons	38
4	Plant growth and physiology	
	<i>Key messages</i>	1
4.1	Germination and emergence issues	2
4.1.1	Germination	2
4.1.2	Emergence.....	3
	<i>Effect of Soil Moisture Content on Seedling Emergence and Early Growth of Some Chickpea Genotypes in WA</i>	4
4.2	Effect of temperature, photoperiod and climate effects on plant growth and physiology	6
4.2.1	Temperature	6
	<i>Cold temperatures</i>	7
	<i>Heat stress</i>	8
4.2.2	Photoperiod.....	9
	<i>Determination of Photoperiod-Sensitive Phase in Chickpea (Cicer arietinum L.)</i>	9
4.2.3	Water and moisture	9
	<i>Growth and yield in chickpea genotypes in response to water stress</i>	10
4.3	Plant growth stages	11
4.3.1	Leaves.....	13
4.3.2	Roots.....	14
4.3.3	Branches	16
4.3.4	Flowering.....	17
4.3.5	Podding.....	18
5	Nutrition and fertiliser	
	<i>Key messages</i>	1
5.1	Nutrient types	2
5.2	Crop removal rates	3
5.2.1	Nutrient budgeting	3
5.3	Identifying nutrient deficiencies	5
5.3.1	Tests for nutrient deficiency	5
5.4	Soil testing	6
5.4.1	Types of test.....	8
	<i>Collecting soil samples for nutrient testing</i>	8

<i>Depth for nutrient sampling</i>	8
<i>Critical values and ranges</i>	9
5.4.2 West Australian Soil Quality Monitoring Program	11
5.5 Plant and/or tissue testing for nutrition levels	11
5.6 Fertiliser	12
5.6.1 Fertiliser toxicity	13
5.7 Nitrogen	14
5.7.1 Chickpeas and nitrogen	16
5.7.2 Deficiency symptoms	16
<i>What to look for</i>	16
5.7.3 Yield potential and nitrogen requirement	18
<i>Nitrogen application to chickpeas is often not worth the cost</i>	19
5.8 Phosphorus	19
5.8.1 Deficiency symptoms	21
<i>What to look for</i>	21
5.8.2 Fate of applied fertiliser	22
5.8.3 Measuring a soil's ability to fix phosphorus	23
5.8.4 Phosphorus retention and removal	23
5.8.5 Leaching and placement of phosphorus	24
5.8.6 Soil P testing	24
5.9 Sulfur	24
5.9.1 Symptoms	25
5.9.2 Applying Sulfur	25
<i>Growth, nitrogen fixation and nutrient uptake by chickpea in response to phosphorus and sulfur application under rainfed conditions in Pakistan</i>	25
5.10 Potassium	26
<i>Diagnosis of potassium deficiency in faba bean and chickpea by plant analysis</i>	26
5.10.1 Symptoms	27
5.10.2 Applying potassium	27
5.11 Micronutrients	28
5.11.1 Zinc	28
<i>Symptoms</i>	28
<i>Applying Zinc</i>	29
<i>Pre-plant treatments</i>	29
<i>Seed treatments</i>	29
<i>Fertilisers applied at sowing</i>	30
<i>Foliar zinc sprays</i>	30
5.11.2 Boron	30
5.11.3 Boron toxicity	31
<i>Boron testing</i>	33
5.11.4 Iron	33
<i>Symptoms</i>	34

5.12 Nutritional deficiencies 34

5.13 Green and Brown Manuring 34

 5.13.1 Outline of procedure..... 35

 5.13.2 Benefits..... 35

 5.13.3 Risks..... 36

 5.13.4 Current level of adoption 36

6 Weed control

Key messages..... 1

 6.1.1 Critical period for weed control..... 2

6.2 Integrated weed management (IWM)..... 3

6.3 Planting control strategies..... 5

 6.3.1 Managing wild oats in chickpeas..... 5

 6.3.2 Row spacing 6

Chemical and Non-chemical Weed Control in Wide Row Lupins and Chickpeas in Western Australia 8

6.4 Herbicides explained 9

 6.4.1 Residual and non-residual 9

 6.4.2 Post-emergent and pre-emergent..... 9

6.5 Mode of Action (MOA) 9

 6.5.1 MOA labelling..... 10

 6.5.2 Grouping by mode of action and ranking by resistance risk 10

 6.5.3 Specific guidelines for Group A herbicides 10

 6.5.4 Specific guidelines for Group B herbicides 11

Broadleaf weed control 12

Grass-weed control..... 12

 6.5.5 Specific guidelines for Group C herbicides 12

 6.5.6 Specific guidelines for Group D herbicides 13

 6.5.7 Specific guidelines for Group F herbicides..... 14

 6.5.8 Specific guidelines for Group I herbicides 14

 6.5.9 Specific guidelines for Group J herbicides 15

 6.5.10 Specific guidelines for Group K herbicides..... 16

 6.5.11 Specific guidelines for Group L herbicides..... 16

No-tillage..... 17

Lucerne..... 17

Horticulture..... 17

 6.5.12 Specific guidelines for Group M herbicides..... 17

 6.5.13 Specific guidelines for Group Z herbicides..... 18

 6.5.14 Herbicide use according to growth stage..... 19

 6.5.15 Getting the best results from herbicides 20

6.6 Summer fallow weed control 20

6.7 Double knock strategies 21

	<i>Getting the best bang for your buck</i>	21
6.8	Pre-emergent herbicides	21
6.8.1	Standard pre-sowing weed control practice—WA	22
6.8.2	Why use pre-emergent herbicides?	22
6.8.3	Herbicide options.....	23
6.8.4	Application.....	25
6.9	Post-plant pre-emergent herbicides	26
	<i>Effect of row spacing, nitrogen and weed control on crop and weed in a wheat—lupin or wheat—chickpea rotation in WA.</i>	28
6.10	In-crop herbicides: knockdowns and residuals	30
6.10.1	Directed sprays.....	31
6.10.2	Shielded sprayers	31
6.11	Conditions for spraying	32
6.11.1	Minimising spray drift.....	33
	<i>Before spraying</i>	33
	<i>During spraying</i>	33
6.11.2	Types of drift.....	34
6.11.3	Factors affecting the risk of spray drift.....	34
	<i>Volatility</i>	35
	<i>Minimising drift</i>	35
	<i>Spray release height</i>	37
	<i>Size of area treated</i>	37
	<i>Capture surface</i>	38
	<i>Weather conditions to avoid</i>	38
6.12	Herbicide tolerance ratings	39
	<i>Herbicide tolerance of new chickpea varieties 2013 trial report, WA</i>	40
6.13	Monitoring	43
6.13.1	Tips for monitoring.....	44
	<i>When to scout, and what to look for in a new paddock or farm</i>	44
6.14	Potential herbicide damage effect	45
	<i>Leaching</i>	45
	<i>Contamination of Spray Equipment</i>	45
	<i>Spray Drift</i>	46
6.14.1	Avoiding herbicide damage	46
6.14.2	Plant-back intervals	47
	<i>Conditions required for breakdown</i>	49
6.15	Herbicide residues	49
6.15.1	Sulfonylurea residues, Group B	50
	<i>Moderate residue levels</i>	50
	<i>Symptoms include:</i>	51
	<i>Highly sensitive crops (in order of susceptibility)</i>	51
	<i>Highly susceptible indicator weeds</i>	51
	<i>Strategy</i>	51

6.15.2 Imidazolinone (imi) residues, Group B.....	51
<i>Moderate residue levels</i>	52
<i>Low residue levels</i>	52
<i>Symptoms include:</i>	52
<i>Highly sensitive crops (in order of susceptibility)</i>	52
<i>Strategy</i>	53
6.15.3 Triazine residues (atrazine), Group C.....	53
<i>Highly susceptible indicator weeds</i>	53
6.15.4 Group I.....	54
6.15.5 Group I residual herbicides.....	55
<i>Strategy</i>	56
6.15.6 Management of herbicide residues in the soil.....	56
6.16 Herbicide resistance.....	57
<i>Annual ryegrass herbicide resistance</i>	59
<i>Glyphosate resistance</i>	59
6.16.1 Practices to minimise herbicide resistance.....	63
6.16.2 WeedSmart farming.....	63
<i>WeedSmart 10-point plan</i>	64
6.16.3 Testing for herbicide resistance.....	64
<i>In-situ testing</i>	64
<i>Herbicide resistance seed tests</i>	65
<i>Syngenta herbicide resistance Quick-Test™</i>	65
6.17 Grazing for weed control.....	65
6.17.1 Grazing stubbles or failed crops.....	67
7 Insect control.....	1
<i>Key messages</i>	1
<i>Insect ID: The Ute Guide</i>	2
7.1.1 Key Integrated pest management (IPM) strategies for chickpeas.....	2
7.2 Pest management process.....	3
7.3 Legal considerations of pesticide use.....	4
7.3.1 Registration.....	4
7.3.2 Labels.....	4
<i>MSDS</i>	4
<i>Permits</i>	4
<i>Always read the label</i>	5
7.4 Native budworm.....	5
<i>Mortality of eggs and caterpillars</i>	8
7.4.1 Varietal resistance or tolerance.....	9
<i>Chickpea, field pea, lentil and faba bean crops:</i>	9
7.4.2 Damage caused by native budworm.....	9
<i>Cost of native budworm</i>	9
7.4.3 Conditions favouring development.....	10
7.4.4 Thresholds for control.....	10

	<i>Threshold tables</i>	10
	<i>Adjusting thresholds</i>	12
7.4.5	Making a decision to control	12
	<i>Selecting control options</i>	13
7.4.6	Management of helicoverpa	13
	<i>Monitoring</i>	13
	<i>Monitoring for adult moths</i>	15
	<i>Recording of monitoring data for decision-making</i>	16
	<i>Chemical control</i>	17
	<i>Biological control</i>	18
7.4.7	Management nearing dessication and harvest	18
7.4.8	Broader management considerations	19
7.4.9	Checking compatibility of products used in mixtures	20
	<i>Compatibility of insecticides with mancozeb formulations</i>	20
7.4.10	Post spray assessments	21
7.5	Aphids	21
	<i>Pulse aphids</i>	21
	<i>Life cycle</i>	23
	<i>Damage</i>	23
	<i>Control</i>	24
7.5.1	Bluegreen aphid	24
	<i>Life cycle</i>	25
	<i>Damage</i>	25
	<i>Control</i>	25
7.5.2	Management of aphids	26
	<i>Monitoring</i>	26
	<i>Chemical control</i>	27
	<i>Biological control</i>	28
	<i>Cultural control</i>	28
7.5.3	Aphids and virus incidence	28
7.5.4	Integrated pest management and viruses	29
7.6	Red legged earth mite (RLEM)	30
7.6.1	Symptoms	30
	<i>What to look for</i>	30
7.6.2	Damage caused by RLEM	32
	<i>RLEM effects:</i>	32
7.6.3	Conditions favouring development	33
7.6.4	Management of RLEM	33
	<i>Monitoring</i>	34
	<i>Chemical control</i>	34
	<i>Biological control</i>	34
	<i>Cultural control</i>	35
7.6.5	RLEM insecticide resistance in WA	35
	<i>How does resistance occur?</i>	35

	<i>Chemical control options</i>	36
	<i>How long does resistance last?</i>	36
	<i>Spread of resistance</i>	36
	<i>Managing resistance</i>	36
	<i>What you can do this season</i>	37
7.7	Lucerne flea	37
7.7.1	Symptoms	38
	<i>What to look for</i>	38
7.7.2	Damage caused by Lucerne flea.....	39
7.7.3	Thresholds for control.....	39
7.7.4	Conditions favouring development.....	39
7.7.5	Management of Lucerne flea.....	40
	<i>Monitoring</i>	40
	<i>Chemical control</i>	40
	<i>Biological control</i>	41
7.8	Cutworms	42
7.8.1	Symptoms	42
	<i>What to look for</i>	42
7.8.2	Damage caused by cutworms.....	44
	<i>Economic and financial considerations</i>	44
7.8.3	Conditions favouring development.....	45
7.8.4	Thresholds for control.....	45
7.8.5	Control	45
	<i>Monitoring</i>	45
	<i>Chemical control</i>	45
	<i>Cultural control</i>	45
	<i>Biological control</i>	46
7.9	Locusts	46
7.9.1	Effect on growing crops	46
7.9.2	Locusts can impact on pulse deliveries	47
	<i>Key points</i> :	47
7.9.3	Management of locusts.....	47
	<i>Control</i>	48
	<i>Spur-throated locust: insecticide spraying guide—WA</i>	48
7.10	Slugs and snails	49
7.10.1	Increase in WA.....	49
	<i>Distribution of slugs and snails</i>	50
7.10.2	Damage caused by slugs and snails	50
7.10.3	Thresholds for control.....	50
7.10.4	Management of slugs and snails	50
	<i>Chemical control</i>	51
	<i>Biological control</i>	52
	<i>Cultural control</i>	52

7.10.5 Monitoring.....	53
<i>How to find slugs</i>	53
<i>How to find snails</i>	53
8 Nematode management	
<i>Key messages</i>	1
8.1 Root-lesion nematode (RLN)	2
<i>Key points</i>	2
8.1.1 <i>Pratylenchus quasitereoides</i> (formerly <i>teres</i>)—WA's home grown RLN.....	4
8.1.2 Varietal resistance or tolerance	5
8.1.3 Damage caused by RLN.....	6
8.1.4 Symptoms	6
<i>Root damage—dark lesions and poor root structure</i>	7
<i>Plant tops—stunted, yellow lower leaves, wilting</i>	8
8.1.5 Conditions favouring development.....	8
8.1.6 Thresholds for control.....	9
8.1.7 Management of RLN	9
<i>Monitoring</i>	10
<i>Soil testing</i>	11
<i>Strategies</i>	12
<i>Nematicides</i>	13
<i>Varietal choice and crop rotation options</i>	13
<i>Yield response in chickpea cultivars and wheat following crop rotations affecting population densities of Pratylenchus thornei and arbuscular mycorrhizal fungi</i>	13
<i>Fallow</i>	14
8.1.8 Breeding resistance	14
<i>Hybridisation of Australian chickpea cultivars with wild Cicer spp. increases resistance to root-lesion nematodes (Pratylenchus thornei and P. neglectus)</i>	14
<i>Highly heritable resistance to root-lesion nematode (Pratylenchus thornei) in Australian chickpea germplasm observed using an optimised glasshouse method and multi-environment trial analysis</i>	15
8.2 Nematodes and crown rot.....	16
8.2.1 Management	16
9 Diseases	
<i>Key messages</i>	1
9.1 Key disease management strategies for chickpea.....	4
9.2 Fungal disease management strategies.....	5
<i>Key strategies</i>	6
9.3 Integrated disease management	7
9.4 Risk assessment	8
9.4.1 Steps in risk assessment.....	8
9.4.2 Paddock selection	8
9.4.3 Regular crop monitoring.....	9

	<i>For Ascochyta blight</i>	9
	<i>For Botrytis grey mould</i>	9
9.4.4	Services and resources available to assist with disease forecasts, disease occurrence and identification.....	10
	<i>Crop disease forecasts</i>	10
	<i>PestFax</i>	10
	<i>Diagnosis tools and services</i>	11
9.5	Ascochyta blight	11
	<i>Economic importance</i>	12
9.5.1	Varietal resistance or tolerance	12
9.5.2	Damage caused by disease.....	12
9.5.3	Symptoms	13
9.5.4	Conditions favouring development.....	15
9.5.5	Management of disease.....	16
	<i>Monitoring</i>	16
	<i>Hygiene</i>	17
	<i>Control</i>	17
9.5.6	Ascochyta blight management in Kabuli	18
	<i>Paddock selection</i>	18
	<i>Seed</i>	18
	<i>Fungicide timing</i>	18
9.5.7	Foliar fungicide programs.....	19
	<i>Economic chickpea production for southern Australia through improved cultivars and strategic management to control Ascochyta blight</i>	19
	<i>Management options for minimizing the damage by Ascochyta blight in chickpea</i>	21
9.6	Botrytis grey mould	22
	<i>Economic importance</i>	22
9.6.1	Varietal resistance or tolerance	22
9.6.2	Damage caused by BGM	22
9.6.3	Symptoms	22
9.6.4	Conditions favouring development.....	25
9.6.5	Management of BGM	25
	<i>Stubble management</i>	25
	<i>Volunteer control (the green bridge)</i>	26
	<i>Seed source and treatment</i>	26
	<i>Seedling emergence</i>	26
	<i>Paddock selection</i>	26
	<i>Sowing time and row spacing</i>	26
	<i>Varietal resistance</i>	27
	<i>Fungicide treatment</i>	27
9.7	Sclerotinia	27
9.7.1	Varietal resistance or tolerance	28
9.7.2	Damage caused by disease.....	28
9.7.3	Symptoms	28

9.7.4	Conditions favouring development.....	29
9.7.5	Management of sclerotinia.....	30
	<i>Before sowing</i>	30
9.8	Phytophthora root rot.....	31
9.8.1	Varietal resistance or tolerance	31
9.8.2	Damage caused by PRR.....	31
9.8.3	Symptoms	31
9.8.4	Phytophthora and waterlogging	33
9.8.5	Conditions favouring development.....	34
9.8.6	Management of PRR.....	34
9.9	Root rots including Damping off (<i>Fusarium</i>, <i>Rhizoctonia</i> and <i>Pythium spp.</i>)	35
9.9.1	Economic importance	35
9.9.2	Symptoms	35
9.9.3	Management options	36
9.10	Collar rot (<i>Sclerotium rolfsii</i>).....	36
9.10.1	Economic importance	36
9.10.2	Symptoms	36
9.10.3	Conditions favouring development.....	37
9.10.4	Management options.....	37
9.11	Fungal disease control.....	37
9.11.1	When to spray.....	37
9.11.2	Principles of spraying.....	38
9.12	Viruses.....	40
	<i>Key points</i>	40
9.12.1	Symptoms	41
9.12.2	Conditions favouring development.....	41
	<i>Types of transmission</i>	42
9.12.3	Management of viruses.....	43
	<i>Better agronomy—better chickpeas</i>	43
	<i>Row spacing and incidence of plants with virus symptoms</i>	45
	<i>Stubble management and incidence of plants with virus symptoms</i>	45
	<i>Non-persistently transmitted viruses</i>	46
	<i>Persistently transmitted viruses</i>	46
10	Plant growth regulators and canopy management	
11	Crop desiccation/spray out	
	<i>Key messages</i>	1
11.1.1	Benefits of desiccation	1
11.2	Croptop, desiccate, harvest or manure?.....	2
11.2.1	When weeds are not the priority	2
	<i>Option 1</i>	2
	<i>Option 2</i>	2

11.2.2	When weeds are the priority, particularly if herbicide resistance exists	2
	<i>Option 1</i>	2
	<i>Option 2</i>	2
11.3	Timing of desiccation	3
	<i>Monitoring for desiccation timing</i>	3
	<i>Seed and pod development</i>	5
	<i>Effect of desiccants on immature seeds</i>	6
11.3.1	Products for the desiccation of chickpea	6
11.4	Croptopping	8
12	Harvest	1
	<i>Key messages</i>	1
12.1	Windrowing	2
12.2	Harvest timing.....	3
12.2.1	Major losses from late harvest include:.....	5
	<i>Loss of yield</i>	5
	<i>Loss of quality</i>	5
	<i>Increased disease and insect risk to pods and seed</i>	6
	<i>Lost marketing opportunities</i>	6
12.2.2	Planning for early harvest	7
12.3	Header modifications and settings.....	8
12.3.1	Draper Fronts	9
12.3.2	Preferred Air Front Setups.....	9
12.3.3	Conventional Headers.....	10
12.3.4	Sieves.....	10
12.3.5	Header Speeds.....	10
12.4	Getting a clean sample.....	11
12.4.1	Perforated screens	11
12.4.2	Harvester speed.....	11
12.4.3	Harvesting in high humidity.....	11
12.4.4	Pick-up fronts	11
12.4.5	Flexible cutter-bar fronts (flexi-fronts)	11
12.4.6	Lodged crops	11
12.5	Fire prevention.....	12
	<i>Harvester fire reduction checklist</i>	12
12.5.1	Harvesting in low-risk conditions.....	13
12.6	Receival standards	14
	<i>Definitions</i>	15
12.7	Harvest weed seed management	15
12.7.1	Factors affecting weed seed removal	16
12.7.2	Strategies for weed seed removal at harvest	17
	<i>Narrow windrow burning</i>	17
	<i>Chaff Carts</i>	18

	<i>Bale direct systems</i>	18
	<i>Integrated Harrington Seed Destructor</i>	19
13	Storage	
	<i>Key messages</i>	1
	<i>Condition of the seed at harvest</i>	2
13.1	How to store chickpeas on-farm	2
	<i>Key points</i>	2
13.1.1	Handling and storage of chickpea seed	3
	<i>Planting-seed selection</i>	3
	<i>Handling</i>	3
	<i>Seed longevity in storage</i>	3
	<i>Moisture and temperature</i>	4
	<i>Weathering damage hinders storage</i>	4
13.1.2	On-farm storage	5
	<i>Established strategies</i>	7
	<i>New products and equipment</i>	8
13.1.3	Silos	8
	<i>Pressure testing</i>	10
	<i>The importance of a gas-tight silo</i>	11
13.1.4	Grain bags	12
	<i>Risks with chickpeas</i>	13
13.1.5	Grain storage—get the economics right	14
	<i>Comparing on-farm grain storage</i>	14
	<i>Summary</i>	15
13.1.6	Hygiene	19
	<i>Where to clean</i>	19
	<i>When to clean</i>	20
	<i>How to clean</i>	20
13.1.7	Aeration cooling	22
	<i>Using aeration cooling for stored grain in Western Australia</i>	22
	<i>Air movement within the stack</i>	24
	<i>The cooling process</i>	24
	<i>Achieving reliable results with aeration cooling</i>	25
	<i>The risks of getting it wrong</i>	25
	<i>Installation and management tips</i>	26
	<i>Monitoring is a must</i>	26
13.1.8	Aeration drying	27
	<i>Management strategies</i>	28
	<i>High airflow for drying</i>	28
	<i>Ducting for drying</i>	29
	<i>Venting for drying</i>	29
	<i>Weather conditions for drying</i>	29
	<i>Phase one of drying</i>	30
	<i>Phase two of drying</i>	30
	<i>Supplementary heating</i>	30

Cooling after drying.....	30
13.1.9 Cooling or drying: making a choice	31
13.1.10 Aeration controllers.....	31
13.1.11 Structural treatments for chickpea storage.....	32
Application.....	33
Silo application.....	33
13.1.12 Fumigation	34
Maximum residue limits.....	35
Phosphine application.....	36
Non-chemical treatment options.....	37
13.2 Monitoring stored chickpeas.....	38
13.3 Grain protectants for storage.....	39
14 Environmental issues	
Key messages.....	1
14.1 Frost issues for chickpeas	1
14.1.1 Impacts on chickpea	3
Damage to vegetative growth.....	3
Damage to flowers and pods	4
14.1.2 Industry costs	5
14.1.3 Managing to lower frost risk.....	6
Problem areas and timings.....	6
Crop and sowing time	7
Spread the risk.....	7
Reduce frost damage	7
Importance of soil moisture	10
14.1.4 Managing frost affected crops.....	11
14.2 Temperature	12
14.2.1 Impact of freezing range (below -1.5°C)	13
14.2.2 Impact of chilling range (-1.5°C–15°C)	13
Tolerance to low temperature.....	16
Response of chickpea genotypes to low-temperature stress during reproductive development.....	18
14.2.3 Heat stress.....	21
Heat-stress-induced reproductive failures in chickpea are associated with impaired sucrose metabolism in leaves and anthers.....	22
14.2.4 Heat and water stress.....	23
Response of chickpea to short periods of high temperature and water stress at different developmental stages	23
14.3 Waterlogging/flooding issues for chickpeas.....	24
14.3.1 Symptoms	25
Effect of waterlogging on stomatal conductance and photosynthesis	26
Effects of waterlogging on chickpeas: influence of timing of waterlogging.	27
14.3.2 Management options for waterlogging.....	27
Identifying problem areas	28

	<i>Raised bed scan overcome waterlogging.....</i>	28
14.4	Drought stress	28
	<i>Response of chickpea to terminal drought: leaf stomatal conductance, pod abscisic acid concentration, and seed set, in WA</i>	29
	<i>Physiological responses of chickpea genotypes to terminal drought in a Mediterranean-type environment, WA.....</i>	30
14.4.1	Managing for drought.....	31
	<i>Foliar nitrogen applications increase the seed yield and protein content in chickpea subject to terminal drought.....</i>	31
14.4.2	Adaptation to drought stress	32
	<i>The role of phenology in adaptation of chickpea to drought.....</i>	33
	<i>GRDC Project ICA00008—Breeding chickpea for drought tolerance and disease resistance.....</i>	34
14.5	Other environmental issues	34
14.5.1	Salinity.....	34
	<i>Effects of salt stress on growth, nodulation, and nitrogen and carbon fixation of ten genetically diverse lines of chickpea</i>	37
14.5.2	Soil chloride levels.....	37
	<i>Agronomic practices and crop choice.....</i>	38
14.5.3	Soil pH.....	39
	<i>Acidic soils.....</i>	39
	<i>Soil pH in WA.....</i>	39
	<i>Alkaline soils.....</i>	40
	<i>Managing soil acidity.....</i>	41
14.5.4	Sodicity	43
	<i>Sodic soils in WA.....</i>	44
	<i>Managing sodic soils.....</i>	45
15	Marketing	
15.1	Selling principles	2
15.1.1	Be prepared.....	2
	<i>When to sell.....</i>	2
	<i>How to sell.....</i>	2
15.1.2	Establish the business risk profile	3
	<i>Production risk profile of the farm</i>	4
	<i>Establishing a target price</i>	5
	<i>Income requirements.....</i>	6
15.1.3	Managing your price	7
	<i>Methods of price management.....</i>	8
15.1.4	Ensuring access to markets	11
	<i>Storage and logistics</i>	11
	<i>Cost of carrying grain</i>	12
15.1.5	Converting tonnes into cash.....	13
	<i>Set up the toolbox.....</i>	13
	<i>How to sell for cash.....</i>	14
	<i>Counterparty risk.....</i>	17

<i>Relative values</i>	17
<i>Contract allocation</i>	19
<i>Read market signals</i>	19
15.2 Western chickpeas: market dynamics and execution	20
15.2.1 Price determinants for western chickpeas.....	20
15.2.2 Ensuring market access for western chickpeas	22
15.2.3 Converting tonnes into cash for western chickpeas	23
16 Current and past research	
17 References	